



## Background



## Methodology



## References

Müller, R.D., Seton, M., Zahirovic, S., Williams, S.E., Matthews, K.J., Wright, N.M., Shephard, G.E., Maloney, K.T., Barnett-Moore, N., Hosseinpour, M., Bower, D.J., Cannon, J., 2016. Ocean Basin Evolution and Global-Scale Plate Reorganization Events Since Pangea Breakup. *Annual Review of Earth and Planetary Sciences* 44, 107-138.


Van Der Meer, D.G., Torsvik, T.H., Spakman, W., Van Hinsbergen, D.J.J., and Amaru, M.L., 2012. Intra-Panthalassa Ocean subduction zones revealed by fossil arcs and mantle structure: *Nature Geoscience*, v. 5, p. 215–219, doi: 10.1038/ngeo1401

Wu, J., J. Suppe, R. Lu, and R. Kanda (2016), Philippine Sea and East Asian plate tectonics since 52 Ma constrained by new subducted slab reconstruction methods, *J. Geophys. Res. Solid Earth*, 121, 4670–4741, doi:10.1002/2016JB012923.

## Acknowledgments

The author thanks Spencer Fuston for technical assistance, the Summer Undergraduate Research Fellowship from the University of Houston's Honor college, and open-source software Gplates. Gocad educational licenses were provided by Paradigm.

## Results: Earth's slab graveyard under the Pacific




Slab Number	Median depth km	0 MA	25 MA	50 MA	75 MA	100 MA	125 MA	150 MA	175 MA	200 MA	225 MA	250 MA
2	1,067											
19	1,132											
3	1,245											
12	1,288											
13	1,421											
1	1,511											
16	1,533											
14	1,687											
4	1,752											
20	1,932											
5	2,086											
17	2,108											
8	2,350											
18	2,351											
15	2,462											
7	2,483											
9	2,483											
6	2,505											
11	2,726											

Plate is more than or equal to 75% visible (i.e. no overlaps) within a plate reconstruction\*

Plate is less than 75% visible (i.e. nearly vanished from Earth's surface)

- Slabs are numbered and median depth was calculated
- The slabs were then compared with time dependent files of accepted plate movement
- The plate was tested on a pass or fail test based on whether 75 percent of its body was visible
- Before 100 million years the predicted plates were viable
- At around 100 million years and after the plates were somehow subducted under the expanding Pacific, and other oceanic basins and continents.



## Conclusions

- These plates have a wide array of depths that primarily correspond to long periods of subduction in the paleo-Pacific region prior to 100 million years ago
- The stark change in visibility across the 100 million year time mark indicates a possible increase in plate tectonic activity (i.e. a plate reorganization) in the paleo-Pacific region